arXiv:1212.2937v1 [math.CO] 12 Dec 2012

Orbit graphs and face-transitivity of k-orbit polytopes

Gabe Cunningham
University of Massachusetts Boston
Boston, Massachusetts, 02125 USA

December 13, 2012

Abstract

The orbit graph of a k-orbit polytope is a graph on k nodes that shows how the flag orbits are related by flag adjacency. Using orbit graphs, we classify k-orbit polytopes and determine when a k-orbit polytope is i-transitive. We then provide an explicit classification of three-orbit polytopes, and we describe a generating set for their automorphism groups.

Key Words: abstract polytope, 2-orbit polytope, k-orbit polytope.

AMS Subject Classification (2000): Primary: 52B15. Secondary: 51M20, 05C25.

1 Introduction

Abstract polytopes are combinatorial generalizations of convex polytopes, maps on surfaces, and tessellations of Euclidean and hyperbolic spaces. A polytope's automorphism group acts naturally on the flags of that polytope, and a *k-orbit polytope* is one where there are k flag orbits. The 1-orbit polytopes, usually called *regular polytopes*, are the most studied, and [5] is the standard reference. Chiral polytopes, an important subclass of the 2-orbit polytopes, have also been studied extensively, though many questions still remain unanswered (see [9] for a thorough survey). More recently, Hubard classified and described the automorphism groups of the 2-orbit polytopes in [3]. A number of interesting results are also known for polytopes with no preassigned number of flag orbits; see [2, 6, 7] for some recent examples.

In the general case of k-orbit polytopes, very little is known. The edge-transitive 4-orbit polyhedra were studied in [4], and an algebraic classification of k-orbit maps (which are essentially abstract polyhedra) is given in [8]. Hypermaps with multiple orbits were studied in [1]. In higher rank, however, no systematic study of k-orbit polytopes has been done.

The key idea used to classify 2-orbit polytopes in [3] is to study the local configuration of flags. That is, given a flag in one orbit, we want to determine which nearby flags are in which orbits. In this paper, we formalize this idea by introducing the *orbit graph* of a polytope. If $\mathcal{P}$ is a k-orbit n-polytope, its orbit graph is a multigraph on k nodes, without loops, and where each arc is labeled with an integer from $\{0, \dots, n-1\}$. Whenever an arc labeled i connects two nodes, this indicates that for any flag Φ in either flag orbit, the flag Φ^i (which differs from Φ only in its i-face) lies in the other orbit. We can classify k-orbit polytopes using their orbit graphs, and so we start in Section 3 by determining some of the necessary properties of an orbit graph.

A number of other properties of $\mathcal{P}$ are apparent from $G(\mathcal{P})$. Corollary 4.3 shows how we can determine the number of orbits $\Gamma(\mathcal{P})$ has on the i-faces using a subgraph $G_i(\mathcal{P})$ of $G(\mathcal{P})$. Similarly, the graph $G_i(\mathcal{P})$ is related to the orbit graphs of the i-faces of $\mathcal{P}$, and Proposition 4.4 and Theorem 4.7 explore this relation. In particular, Theorem 4.7 generalizes the well-known result that the $(n-2)$-faces of a chiral polytope are always regular [10].

In general, there are many distinct orbit graphs for k-orbit n-polytopes, making explicit analysis infeasible. When $k = 3$, however, there are only $2n-3$ different orbit graphs, divided into two families. In Section 5, we apply the results of the previous sections to three-orbit polytopes. We then provide a generating set for the automorphism group of each class of three-orbit polytope.

2 Background

For general background information on abstract polytopes, the reader is directed to [5, Chs. 2, 3]. Here we will review the essential concepts.

Let $\mathcal{P}$ be a ranked partially ordered set whose elements will be called *faces*. The faces of $\mathcal{P}$ will range in rank from -1 to n, and a face of rank j is called a *j-face*. The 0-faces, 1-faces, and $(n-1)$-faces are also called *vertices*, *edges*, and *facets*, respectively. A *flag* of $\mathcal{P}$ is a maximal chain. We say that two flags are *adjacent* if they differ in exactly one face, and that they are *j-adjacent* if they differ only in their j-face. If F and G are faces of $\mathcal{P}$ such that $F \leq G$, then the *section* G/F consists of those faces H such that $F \leq H \leq G$.

We say that $\mathcal{P}$ is an *(abstract) polytope of rank n*, also called an *n-polytope* (or a *polyhedron* if $n = 3$), if it satisfies the following four properties:

(a) There is a unique greatest face F_n of rank n and a unique least face F_{-1} of rank -1.

(b) Each flag of $\mathcal{P}$ has $n+2$ faces.

(c) $\mathcal{P}$ is *strongly flag-connected*, meaning that if Φ and Ψ are two flags of $\mathcal{P}$, then there is a sequence of flags $\Phi = \Phi_0, \Phi_1, \dots, \Phi_k = \Psi$ such that for $i = 0, \dots, k-1$, the flags Φ_i and Φ_{i+1} are adjacent, and each Φ_i contains $\Phi \cap \Psi$.

(d) (Diamond condition): Whenever $F < G$, where F is a $(j-1)$-face and G is a $(j+1)$-face for some j, then there are exactly two j-faces H with $F < H < G$.

Note that due to the diamond condition, any flag Φ has a unique j-adjacent flag (denoted Φ^j) for each $j = 0, 1, \dots, n-1$. Furthermore, if $|i-j| \geq 2$, then $\Phi^{i,j} = \Phi^{j,i}$.

If F is a j-face and G is a k-face of a polytope with $F \leq G$, then the section G/F is a $(k-j-1)$-polytope itself. We can identify a face F with the section F/F_{-1}, since if F is a

j-face, then F/F_{-1} is a j-polytope. We call the section F_n/F the *co-face at* F; the co-face at a vertex is also called a *vertex-figure.*

For polytopes $\mathcal{P}$ and $\mathcal{Q}$, an *isomorphism* from $\mathcal{P}$ to $\mathcal{Q}$ is an incidence- and rank-preserving bijection on the set of faces. An isomorphism from $\mathcal{P}$ to itself is an *automorphism* of $\mathcal{P}$, and the group of all automorphisms of $\mathcal{P}$ is denoted $\Gamma(\mathcal{P})$. If φ is an automorphism of $\mathcal{P}$ and Φ is a flag of $\mathcal{P}$, then $\Phi^i\varphi = (\Phi\varphi)^i$, so that the action of any automorphism is completely determined by the action on a single flag.

There is a natural action of $\Gamma(\mathcal{P})$ on the flags of $\mathcal{P}$. In fact, this action is semiregular; that is, the only automorphism with any fixed points is the identity. As a result, the flag orbits of $\mathcal{P}$ all have the same size, and $\Gamma(\mathcal{P})$ acts regularly on each orbit. If there are k flag orbits under the natural action of $\Gamma(\mathcal{P})$, we say that $\mathcal{P}$ is a *k-orbit polytope.* If there is a single flag orbit, so that the action of $\Gamma(\mathcal{P})$ is transitive on the flags, then we say that $\mathcal{P}$ is a *regular polytope.*

2-orbit polytopes were classified in [3] as follows. If $\mathcal{P}$ is a 2-orbit n-polytope and the flags Φ and Φ^i are in the same orbit (for some $i \in \{0, \ldots, n-1\}$) then every flag Ψ lies in the same orbit as its i-adjacent flag Ψ^i. Therefore, we can classify the 2-orbit polytopes by a proper subset $I \subsetneq \{0, \ldots, n-1\}$. We say that $\mathcal{P}$ is in *class* 2_I if, for any flag Φ, the flags Φ and Φ^i are in the same orbit precisely when $i \in I$. Thus, there are $2^n - 1$ classes of 2-orbit n-polytopes. We usually write $2_{i_1,\ldots,i_k}$ rather than $2_{\{i_1,\ldots,i_k\}}$, so that in rank 3, the classes are denoted by 2, 2_0, 2_1, 2_2, $2_{0,1}$, $2_{0,2}$, and $2_{1,2}$. 2-orbit n-polytopes in class 2 are also called *chiral polytopes* (see [10]).

3 Orbit graphs

In studying the structure of a k-orbit polytope $\mathcal{P}$, we are naturally led to the idea of studying the local flag configuration of $\mathcal{P}$. In other words, starting from a flag Φ, we want to determine which flags near Φ are in which flag orbits. If Ψ is in the same orbit as Φ, then there is an automorphism $\varphi \in \Gamma(\mathcal{P})$ such that $\Phi\varphi = \Psi$. Then

$$\Phi^i\varphi = (\Phi\varphi)^i = \Psi^i,$$

so that Φ^i and Ψ^i are also in the same orbit. As a result, there is a natural way to associate to $\mathcal{P}$ a graph called the *orbit graph* of $\mathcal{P}$, denoted $G(\mathcal{P})$, as follows. The graph $G(\mathcal{P})$ has k nodes, with each one corresponding to a flag orbit. An arc labeled i connects one node to another if, whenever Φ is a flag in one of the corresponding orbits, then Φ^i is a flag in the other orbit. In general, then, two nodes may be connected by multiple arcs. By convention, we omit all loops, which would occur when Φ and Φ^i lie in the same orbit.

Since Φ and Ψ are in the same orbit if and only if Φ^i and Ψ^i are in the same orbit, each node of $G(\mathcal{P})$ is incident on at most one arc with a given label. In other words, the arc labels yield a proper arc coloring of $G(\mathcal{P})$. Furthermore, if $\mathcal{P}$ is an n-polytope, then each node of $G(\mathcal{P})$ has degree n or less.

We start by collecting some simple properties of orbit graphs.

Proposition 3.1. *The orbit graph of a polytope is a connected graph.*

Proof. Since polytopes are flag-connected, it is possible to get from any flag to any other through a sequence of flag adjacencies. Therefore, any connected component of the orbit graph must contain all of the flags. Then since each flag is in a single orbit, the orbit graph must be connected. $\square$

Note that if $\mathcal{P}$ is a 2-orbit n-polytope in class 2_I, where $I \subset \{0, \ldots, n-1\}$, then the set of arc labels that appear in $G(\mathcal{P})$ is $\{0, \ldots, n-1\} \setminus I$. Since I must be a proper subset of $\{0, \ldots, n-1\}$, we see in another way that there is at least one arc connecting the nodes of $G(\mathcal{P})$.

Proposition 3.2. *Suppose that $\mathcal{P}$ is a k-orbit n-polytope and that $0 \leq i < j \leq n-1$ with $i < j-1$. Then the subgraph of $G(\mathcal{P})$ induced by the arcs with label i and j is a disjoint union of single edges, 2-cycles, and 4-cycles.*

Proof. If $i < j-1$, then $\Phi = \Phi^{i,j,i,j}$ for any flag Φ. Therefore, starting from any node in $G(\mathcal{P})$ and moving along arcs labeled i, j, i, and j (in that order), we must get back to where we started. The result follows. $\square$

Proposition 3.2 puts many strong restrictions on the structure of $G(\mathcal{P})$, as we shall see with the next several results.

Proposition 3.3. *If $G(\mathcal{P})$ has no 4-cycles, then it has no 3-cycles.*

Proof. Suppose $G(\mathcal{P})$ has a 3-cycle. Then all three arcs of this cycle must have different labels, since the labels form a proper arc-coloring. In particular, if i is the smallest label appearing in the cycle and j is the largest, $i < j-1$. Therefore, in the subgraph of $G(\mathcal{P})$ induced by arcs with label i and j, these two arcs share one endpoint but not the other. Then Proposition 3.2 says that these arcs must be part of a 4-cycle, which proves the result. $\square$

Corollary 3.4. *If $\mathcal{P}$ is a three-orbit polytope, its orbit graph has no 3-cycles.*

Proposition 3.5. *Let $\mathcal{P}$ be a k-orbit polytope with $k \geq 3$. Suppose that there are nodes u and v in $G(\mathcal{P})$ such that removing all arcs between them disconnects $G(\mathcal{P})$, and let i be one of the labels appearing on arcs between u and v. Then any arc that is incident to u or to v but not both has label $i-1$ or $i+1$.*

Proof. Let i be one of the labels appearing on arcs between u and v. Suppose that there is an arc that is incident on u and $w \neq v$, and that this arc is labeled j, with $|j-i| \geq 2$. Then in the subgraph of $G(\mathcal{P})$ induced by arcs with label i and j, the nodes u, v, and w all lie in the same connected component. Then by Proposition 3.2, there is a 4-cycle that contains u and v. But then u and v remain connected even if we remove all arcs between them, contradicting our assumption. The same argument works if we consider an arc that is incident on v and $x \neq u$, and the result follows. $\square$

Corollary 3.6. *Let $\mathcal{P}$ be a k-orbit polytope with $k \geq 3$. Suppose that there are nodes u and v in $G(\mathcal{P})$ such that removing all arcs between them disconnects $G(\mathcal{P})$. Then*

(a) *There are at most two arcs connecting u to v.*

(b) *There are at most three arcs incident on u and at most three arcs incident on v.*

Proof. Since $k \geq 3$ and $G(\mathcal{P})$ is connected, there is some arc that is incident on one of the nodes u or v but not the other. Without loss of generality, assume there is an arc labeled j that is incident on u and some other node $w \neq v$. Proposition 3.5 says that the only possible labels for arcs connecting u to v are $j+1$ and $j-1$, so there are at most two arcs connecting u to v. Furthermore, if there are two such arcs, then by Proposition 3.5, any other arc incident to u must have label j, and therefore the only arcs incident to u are the three we have mentioned. On the other hand, if there is only a single arc connecting u to v with the label i, then any other arc that is incident to u but not v has label $i \pm 1$. Therefore, there are at most two arcs that are incident to u and not v, yielding three arcs total that are incident to u. The result then follows by symmetry. ☐

We have cut down considerably the number of possible orbit graphs, but many possibilities still remain, even for modest values of k and n. For example, Figure 1 shows the 22 orbit graphs for 4-orbit polyhedra that satisfy all of the conditions in this section (and these correspond to the 22 classes of 4-orbit maps in [8, Table 1]). We will see in Section 5, however, that the orbit graphs of three-orbit polytopes always fall into one of two families, making explicit analysis possible.

4 Face transitivity

An n-polytope $\mathcal{P}$ is said to be *i-transitive* if $\Gamma(\mathcal{P})$ acts transitively on the set of i-faces of $\mathcal{P}$, and *fully transitive* if it is i-transitive for each $0 \leq i \leq n-1$. In [3], Hubard proves that 2-orbit n-polytopes in class 2_I are fully transitive if $|I| < n-1$, and they are i-transitive for $i \in I$ if $|I| = n-1$. The arguments she uses in Lemma 3 and Theorem 5 can be generalized to k-orbit polytopes by using orbit graphs. Let us define $G_i(\mathcal{P})$ to be the full subgraph of $G(\mathcal{P})$ obtained by deleting all arcs with label i.

Lemma 4.1. *Let Φ and Ψ be flags of a k-orbit n-polytope $\mathcal{P}$. If Φ and Ψ have the same i-face, then they lie in orbits in the same connected component of $G_i(\mathcal{P})$.*

Proof. Let Φ and Ψ be flags with the same i-face F. By the strong flag-connectivity of polytopes, there is a sequence of flags

$$\Phi = \Phi_0, \Phi_1, \ldots, \Phi_m = \Psi$$

such that each Φ_j is adjacent to Φ_{j+1} and such that each Φ_j contains $\Phi \cap \Psi$. Since Φ and Ψ both contain the i-face F, it follows that each flag Φ_j contains that same i-face. Therefore, for each j, $\Phi_{j+1} \neq \Phi_j^i$. Then it follows that we can get from the orbit of Φ to the orbit of Ψ in $G(\mathcal{P})$ without using any arcs labeled i, and so these orbits must lie in the same connected component of $G_i(\mathcal{P})$. ☐



Figure 1: All possible orbit graphs for 4-orbit polyhedra

As a result of Lemma 4.1, we can associate a single connected component of $G_i(\mathcal{P})$ to each i-face of $\mathcal{P}$. Furthermore, the connected components of $G_i(\mathcal{P})$ correspond to the orbits of the i-faces of $\mathcal{P}$:

Theorem 4.2. *Let $\mathcal{P}$ be a k-orbit n-polytope, and let F and F' be i-faces of $\mathcal{P}$. Then there is an automorphism $\gamma \in \Gamma(\mathcal{P})$ sending F to F' if and only if F and F' belong to (flags in) the same connected component of $G_i(\mathcal{P})$.*

Proof. First, suppose that such an automorphism γ exists. Let Φ be a flag with i-face F, and let Ψ be a flag with i-face F'. Since γ sends F to F', the flag $\Phi\gamma$ also has i-face F'. Then by Lemma 4.1, the orbits of $\Phi\gamma$ and Ψ lie in the same connected component of $G_i(\mathcal{P})$. Furthermore, Φ and $\Phi\gamma$ lie in the same orbit, since γ is an automorphism, and thus Φ and Ψ lie in the same connected component of $G_i(\mathcal{P})$.

Conversely, suppose that F and F' belong to (flags in) the same connected component of $G_i(\mathcal{P})$. That is, suppose there is a flag Φ with i-face F and a flag Ψ with i-face F' such that Φ and Ψ lie in the same connected component. That means that there is a path in $G_i(\mathcal{P})$ from the orbit of Φ to the orbit of Ψ, and this path induces a sequence of flags

$$\Phi = \Phi_0, \ldots, \Phi_m,$$

where each $\Phi_{j+1} = \Phi_j^t$ for some $t \neq i$, and such that Φ_m lies in the same orbit as Ψ. In particular, each Φ_{j+1} has the same i-face as Φ_j, and thus Φ_m has i-face F. Now, since

Φ_m is in the same orbit as Ψ, there is an automorphism γ that sends Φ_m to Ψ, and this automorphism must then send F to F'. ☐

Corollary 4.3. *The number of orbits of $\Gamma(\mathcal{P})$ on the i-faces of $\mathcal{P}$ is equal to the number of connected components of $G_i(\mathcal{P})$. In particular, $\mathcal{P}$ is i-transitive if and only if $G_i(\mathcal{P})$ is connected.*

Using Corollary 4.3 and looking back at Figure 1, we note that only the last two graphs correspond to fully transitive 4-orbit polyhedra. More generally, we note that the orbit graph of a fully transitive k-orbit polytope must have at least k arcs; if there are only $k-1$ arcs, then one of the subgraphs $G_i(\mathcal{P})$ will have too few edges to remain connected.

Theorem 4.2 says that the connected components of $G_i(\mathcal{P})$ correspond to the different orbits of i-faces. It is natural to wonder whether the orbit graph of (the polytope corresponding to) an i-face is related to its corresponding connected component. We start with the following simple observation:

Proposition 4.4. *Let F be an i-face of the polytope $\mathcal{P}$, let $\mathcal{K} = F/F_{-1}$, and let C be the connected component of $G_i(\mathcal{P})$ that contains flags with the face F. Then each orbit of $\mathcal{K}$ is obtained by identifying one or more of the orbits that belong to C. In particular, $\mathcal{K}$ has at most one flag orbit for each node of C.*

Proof. We start by noting that any flag Φ of $\mathcal{P}$ that contains F naturally induces a flag $\overline{\Phi}$ of $\mathcal{K}$. Now, let Φ and Ψ be flags that lie in the same orbit of $\mathcal{P}$, and suppose that they both contain F. Now, since Φ and Ψ lie in the same orbit, there is some automorphism $\gamma \in \Gamma(\mathcal{P})$ such that $\Phi\gamma = \Psi$. Then γ fixes F, and so it induces an automorphism $\overline{\gamma}$ of $\mathcal{K}$ such that $\overline{\Phi}\overline{\gamma} = \overline{\Psi}$. Therefore, the induced flags $\overline{\Phi}$ and $\overline{\Psi}$ are in the same orbit. Thus, to each orbit O of $\mathcal{P}$ that contains flags with the face F, we can associate a single orbit $\overline{O}$ of $\mathcal{K}$. Since every flag that contains F is part of an orbit in C (by Lemma 4.1), the result follows. ☐

Proposition 4.4 shows that each orbit of $\mathcal{K}$ is obtained by identifying one or more orbits of $\mathcal{P}$. In fact, this identification respects flag adjacency in the following sense:

Theorem 4.5. *Let F be an i-face of the polytope $\mathcal{P}$, let $\mathcal{K} = F/F_{-1}$, and let $j \leq i-1$. Let O_1 and O_2 be flag orbits of $\mathcal{P}$ that contain flags with i-face F, and let $\overline{O_1}$ and $\overline{O_2}$ be the induced flag orbits of $\mathcal{K}$.*

(a) *If there is no arc labeled j incident on O_1 in $G(\mathcal{P})$, then there is no arc labeled j incident on $\overline{O_1}$ in $G(\mathcal{K})$.*

(b) *If $O_1 \neq O_2$ and $\overline{O_1} \neq \overline{O_2}$, and if O_1 and O_2 are connected by an arc labeled j in $G(\mathcal{P})$, then $\overline{O_1}$ and $\overline{O_2}$ are connected by an arc labeled j in $G(\mathcal{K})$.*

(c) *If $O_1 \neq O_2$ and $\overline{O_1} = \overline{O_2}$, and if O_1 and O_2 are connected by an arc labeled j in $G(\mathcal{P})$, then there is no arc labeled j incident on $\overline{O_1}$ in $G(\mathcal{K})$.*

Proof. Let Ψ be a flag of $\mathcal{K}$ in $\overline{O_1}$. Then there is a flag Φ of $\mathcal{P}$ in O_1 such that $\overline{\Phi} = \Psi$; that is, Φ contains F, and Φ and Ψ agree on all faces of rank i or less. Now, since $j \leq i-1$, the flag Φ^j also contains F, and so it induces a flag $\overline{\Phi^j}$ of $\mathcal{K}$. Then

$$\begin{aligned}\overline{\Phi^j} &= \overline{\Phi}^j \\ &= \Psi^j.\end{aligned}$$

Therefore, if Φ^j is in O_1, then Ψ^j is in $\overline{O_1}$, and if Φ^j is in O_2, then Ψ^j is in $\overline{O_2}$. Now, if there is no arc labeled j incident on O_1, then Φ^j is in O_1, Ψ^j is in $\overline{O_1}$, and there is no arc labeled j incident on $\overline{O_1}$ (since we omit loops). On the other hand, if there is an arc labeled j connecting O_1 to O_2, then Φ^j is in O_2 and thus Ψ^j is in $\overline{O_2}$. Therefore, if $\overline{O_1} \neq \overline{O_2}$, there is an arc labeled j that connects them in $G(\mathcal{K})$, and if $\overline{O_1} = \overline{O_2}$, there is no arc labeled j that is incident on $\overline{O_1}$ (since, again, we omit loops). ☐

Corollary 4.6. *Let $\mathcal{P}$ be a k-orbit polytope, let F be one of its i-faces, and let $\mathcal{K} = F/F_{-1}$. If $\mathcal{K}$ is a k-orbit polytope, then $\mathcal{P}$ is i-transitive and $G(\mathcal{K})$ is obtained from $G(\mathcal{P})$ by deleting all arcs with label i or higher.*

Proof. If $\mathcal{K}$ has k orbits, then the connected component C of $G_i(\mathcal{P})$ that contains F must have at least k nodes. Since $G_i(\mathcal{P})$ has only k nodes itself, it follows that C is the whole graph $G_i(\mathcal{P})$. Therefore, $G_i(\mathcal{P})$ is connected, and thus $\mathcal{P}$ is i-transitive, by Corollary 4.3. Furthermore, Theorem 4.5 implies that, for each $j \leq i-1$, the orbit graph $G(\mathcal{K})$ has all of the arcs labeled j that come from $G(\mathcal{P})$. ☐

Let us look at an example to see how we can use Theorem 4.5. Suppose $\mathcal{P}$ is a three-orbit polyhedron with the following orbit graph.



Now, $G_2(\mathcal{P})$ is the whole graph $G(\mathcal{P})$, and so any facet F we choose will have the same connected component $C = G(\mathcal{P})$. Let $\mathcal{K} = F/F_{-1}$. If each orbit of $\mathcal{P}$ corresponds to a unique orbit of $\mathcal{K}$, then $\mathcal{K}$ has 3 orbits, and Corollary 4.6 tells us that $G(\mathcal{K}) \simeq G(\mathcal{P})$. Suppose instead that the first two orbits are identified to a single orbit O in $\mathcal{K}$. Then Theorem 4.5 says that O is not incident on any arcs labeled 1 or 2. Furthermore, since the first two orbits are connected by an arc labeled with 0, the orbit O is not incident on any arcs labeled 0 in $G(\mathcal{K})$. Therefore, O is an isolated vertex of $G(\mathcal{K})$, and by Proposition 3.1, it follows that $G(\mathcal{K})$ consists of a single node. Therefore, $\mathcal{K}$ is regular.

Using similar arguments, we can show that identifying any other pair of orbits leads to the same conclusion. Therefore, the facets of $\mathcal{P}$ are either all regular or all three-orbit polytopes with the same orbit graph as $\mathcal{P}$.

Though Proposition 4.4 and Theorem 4.5 are quite useful, there are some phenomena they fail to explain. For example, the $(n-2)$-faces of a chiral polytope are always regular, whereas all we can conclude from Proposition 4.4 is that the $(n-2)$-faces are either regular or chiral. However, a slight modification of Theorem 4.5 can explain and generalize this property of chiral polytopes.

Theorem 4.7. *Let F be an i-face of the polytope $\mathcal{P}$, let $\mathcal{K} = F/F_{-1}$, and let O_1 and O_2 be distinct orbits of $\mathcal{P}$ that contain flags with the i-face F. Let $\overline{O_1}$ and $\overline{O_2}$ be the induced flag orbits of $\mathcal{K}$. If O_1 and O_2 are connected by an arc labeled j in $G(\mathcal{P})$, where $j > i$, then $\overline{O_1} = \overline{O_2}$.*

Proof. Let Φ be a flag in O_1 that contains the i-face F, let $j > i$, and suppose that O_1 and O_2 are connected by an arc labeled j in $G(\mathcal{P})$. Then Φ^j also contains F and is in the orbit O_2. Furthermore, since $j > i$, Φ and Φ^j induce the same flag of $\mathcal{K}$; that is, $\overline{\Phi} = \overline{\Phi^j}$. Since $\overline{\Phi}$ lies in $\overline{O_1}$ and $\overline{\Phi^j}$ lies in $\overline{O_2}$, it follows that $\overline{O_1} = \overline{O_2}$. ☐

For example, let $\mathcal{P}$ be a chiral n-polytope. Then $G(\mathcal{P})$ consists of two nodes connected by n arcs, one for each label in $\{0, \dots, n-1\}$. Since there is an arc labeled $n-1$, Theorem 4.7 says that we must identify the two orbits in order to get an orbit diagram for any of the $(n-2)$-faces, and thus the $(n-2)$-faces are regular.

5 Three-orbit polytopes

In general, the classification of k-orbit polytopes remains difficult, even with the various restrictions placed on orbit graphs. Nevertheless, the case $k = 3$ proves tractable, and we analyze it here.

Let $\mathcal{P}$ be a three-orbit n-polytope. Corollary 3.4 tells us that $G(\mathcal{P})$ has no cycles, and therefore we can apply Proposition 3.5. As a result, there are only 2 basic families of orbit graph:



and



We shall say that a three-orbit n-polytope is in class $3_{i,i+1}$ if it has the former orbit graph, and in class 3_i if it has the latter. In the first case, i must be between 0 and $n-2$, yielding $n-1$ different classes; in the second case, i must be between 1 and $n-2$, yielding $n-2$ more classes. In each graph, we have colored one of the nodes black to indicate our choice of default orbit for a base flag. We shall refer to the corresponding orbit as the *black orbit*.

By applying Corollary 4.3, we obtain the following result:

Proposition 5.1. *A three-orbit polytope in class $3_{i,i+1}$ is j-transitive if and only if $j \neq i$ and $j \neq i+1$, and a three-orbit polytope in class 3_i is j-transitive if and only if $j \neq i$. In particular, no three-orbit polytope is fully transitive.*

We can also apply Theorem 4.7 to obtain the following:

Proposition 5.2. *If $\mathcal{P}$ is a three-orbit polytope in class $3_{i,i+1}$ or 3_i, then it has regular i-faces.*

Proof. In both cases, the graph $G_i(\mathcal{P})$ consists of an isolated node and two other nodes connected by one or two arcs. The i-faces that correspond to the isolated node have a single induced flag orbit, by Proposition 4.4, and so they are regular. Furthermore, Theorem 4.7 tells us that, since the other two nodes are connected by an arc labeled $i+1$, they must be identified into a single flag orbit of the corresponding i-face, and so those i-faces are regular too. ☐

We now investigate the automorphism group of three-orbit polytopes. Let $\mathcal{P}$ be an n-polytope in class $3_{i,i+1}$, and let Φ be a base flag in the black orbit. Then Φ^j is in the same orbit as Φ for every $j \neq i$, and thus for each such j, there must be an automorphism ρ_j such that $\Phi\rho_j = \Phi^j$. More generally, to any finite sequence $w = (w_1, w_2, \ldots, w_r)$ with $0 \leq w_j \leq n-1$, we can associate a walk in $G(\mathcal{P})$ that starts at the black node. If the associated walk also ends at the black node, then there is an automorphism α_w that sends Φ to $\Phi^{w_1,\ldots,w_r}$.

Theorem 5.3. *Let $\mathcal{P}$ be a three-orbit n-polytope in class $3_{i,i+1}$, and let Φ be a base flag in the black orbit.*

(a) *If $i = 0$, then*
$$\Gamma(\mathcal{P}) = \langle \{\rho_j\}_{j\neq i}, \alpha_{(i,\,i+1,\,i,\,i+1,\,i)}, \alpha_{(i,\,i+1,\,i+2,\,i+1,\,i)} \rangle.$$

(b) *If $1 \leq i \leq n-3$, then*
$$\Gamma(\mathcal{P}) = \langle \{\rho_j\}_{j\neq i}, \alpha_{(i,\,i-1,\,i)}, \alpha_{(i,\,i+1,\,i,\,i+1,\,i)}, \alpha_{(i,\,i+1,\,i+2,\,i+1,\,i)} \rangle.$$

(c) *If $i = n-2$, then*
$$\Gamma(\mathcal{P}) = \langle \{\rho_j\}_{j\neq i}, \alpha_{(i,\,i-1,\,i)}, \alpha_{(i,\,i+1,\,i,\,i+1,\,i)} \rangle.$$

Proof. First, we note that each α_w given really is an automorphism, since in each case, the walk w forms a closed walk if we start at the black node. To prove that these automorphisms generate the group, it suffices to show that we can send Φ to any other flag in the same orbit using only the given automorphisms. Consider another flag Ψ in the same orbit as Φ; since polytopes are flag-connected, we can write $\Psi = \Phi^{w_1,\ldots,w_r}$, and we can assume that $w_1 \neq w_2$. If $r = 1$, then $w_1 \neq i$ (since Ψ is in the same orbit as Φ), and $\Psi = \Phi\rho_{w_1}$. Otherwise, suppose that $r \geq 2$. If we can show that $\Psi = \Psi'\gamma$, where $\Psi' = \Phi^{w'_1,\ldots,w'_{r-1}}$ and γ is one of the given automorphisms, then the claim will follow by induction on r. We consider several cases:

(a) If $w_1 \neq i$, then $\Psi = (\Phi^{w_2,\ldots,w_r})\rho_{w_1}$.

(b) If $w_1 = i$ and $|w_2 - w_1| \geq 2$, then we can switch w_1 and w_2 and reduce to the previous case.

(c) If $w_1 = i$ and $w_2 = i-1$, then $i \geq 1$ and $\Psi = \Phi^{i,i-1,w_3,\ldots,w_r} = (\Phi^{i,w_3,\ldots,w_r})\alpha_{(i,\,i-1,\,i)}$.

(d) If $w_1 = i$, $w_2 = i+1$, and $|w_3 - w_2| \geq 2$, then we can switch w_3 and w_2, which reduces to one of the previous two cases.

(e) If $w_1 = i$, $w_2 = i+1$, and $w_3 = i+2$, then $i \leq n-3$ and $\Psi = \Phi^{i,i+1,i+2,w_4,\ldots,w_r} = (\Phi^{i,i+1,w_4,\ldots,w_r})\alpha_{(i,\,i+1,\,i+2,\,i+1,\,i)}$.

(f) If $w_1 = i$, $w_2 = i+1$, and $w_3 = i$, then $\Psi = \Phi^{i,i+1,i,w_4,\ldots,w_r} = (\Phi^{i,i+1,w_4,\ldots,w_r})\alpha_{(i,\,i+1,\,i,\,i+1,\,i)}$.

Since every sequence $(w_1, \ldots, w_r)$ fits into one of the above cases (except for the singleton sequence (i), which cannot arise), the claim follows. $\square$

A similar analysis is possible for the class 3_i:

Theorem 5.4. *Let $\mathcal{P}$ be a three-orbit n-polytope in class 3_i, and let Φ be a base flag in the black orbit.*

(a) *If $n = 3$, then*

$$\Gamma(\mathcal{P}) = \langle \{\rho_j\}_{j \neq i}, \alpha_{(i,\,i-1,\,i+1,\,i)}, \alpha_{(i,\,i-1,\,i,\,i-1,\,i)}, \alpha_{(i,\,i+1,\,i,\,i+1,\,i)} \rangle.$$

(b) *If $n \geq 4$ and $i = 1$, then*

$$\Gamma(\mathcal{P}) = \langle \{\rho_j\}_{j \neq i}, \alpha_{(i,\,i-1,\,i+1,\,i)}, \alpha_{(i,\,i-1,\,i,\,i-1,\,i)}, \alpha_{(i,\,i+1,\,i,\,i+1,\,i)}, \alpha_{(i,\,i+1,\,i+2,\,i+1,\,i)} \rangle.$$

(c) *If $n \geq 4$ and $2 \leq i \leq n-3$, then*

$$\Gamma(\mathcal{P}) = \langle \{\rho_j\}_{j \neq i}, \alpha_{(i,\,i-1,\,i+1,\,i)}, \alpha_{(i,\,i-1,\,i-2,\,i-1,\,i)}, \alpha_{(i,\,i-1,\,i,\,i-1,\,i)}, \alpha_{(i,\,i+1,\,i,\,i+1,\,i)}, \alpha_{(i,\,i+1,\,i+2,\,i+1,\,i)} \rangle.$$

(d) *If $n \geq 4$ and $i = n-2$, then*

$$\Gamma(\mathcal{P}) = \langle \{\rho_j\}_{j \neq i}, \alpha_{(i,\,i-1,\,i+1,\,i)}, \alpha_{(i,\,i-1,\,i-2,\,i-1,\,i)}, \alpha_{(i,\,i-1,\,i,\,i-1,\,i)}, \alpha_{(i,\,i+1,\,i,\,i+1,\,i)} \rangle.$$

Proof. As before, we note that each α_w given really is an automorphism, since in each case, the walk w forms a closed walk if we start at the black node. To prove that these automorphisms generate the group, it suffices to show that we can send Φ to any other flag in the same orbit using only the given automorphisms. Consider another flag Ψ in the same orbit as Φ; since polytopes are flag-connected, we can write $\Psi = \Phi^{w_1,\ldots,w_r}$, and we can assume that $w_1 \neq w_2$. If $r = 1$, then $w_1 \neq i$ (since Ψ is in the same orbit as Φ), and $\Psi = \Phi\rho_{w_1}$. Otherwise, suppose that $r \geq 2$. If we can show that $\Psi = \Psi'\gamma$, where $\Psi' = \Phi^{w'_1,\ldots,w'_{r-1}}$ and γ is one of the given automorphisms, then the claim will follow by induction on r. We consider several cases:

(a) If $w_1 \neq i$, then $\Psi = (\Phi^{w_2,\ldots,w_r})\rho_{w_1}$.

(b) If $w_1 = i$ and $|w_2 - w_1| \geq 2$, then we can switch w_1 and w_2 and reduce to the previous case.

(c) If $w_1 = i$, $w_2 = i-1$, and $w_3 \leq i-3$ or $w_3 \geq i+2$, then we can switch w_2 and w_3 and reduce to the previous case.

(d) If $w_1 = i$, $w_2 = i-1$, and $w_3 = i+1$, then $\Psi = \Phi^{i,i-1,i+1,w_4,\ldots,w_r} = (\Phi^{i,w_4,\ldots,w_r})\alpha_{(i,\,i-1,\,i+1,\,i)}$.

(e) If $w_1 = i$, $w_2 = i-1$, and $w_3 = i$, then $\Psi = \Phi^{i,i-1,i,w_4,\ldots,w_r} = (\Phi^{i,i-1,w_4,\ldots,w_r})\alpha_{(i,\,i-1,\,i,\,i-1,\,i)}$.

(f) If $w_1 = i$, $w_2 = i-1$, and $w_3 = i-2$, then $i \geq 2$ and $n \geq 4$, and $\Psi = \Phi^{i,i-1,i-2,w_4,\ldots,w_r} = (\Phi^{i,i-1,w_4,\ldots,w_r})\alpha_{(i,\,i-1,\,i-2,\,i-1,\,i)}$.

(g) If $w_1 = i$, $w_2 = i+1$, and $w_3 \leq i-1$ or $w_3 \geq i+3$, then we can switch w_2 and w_3 and reduce to one of the previous cases.

(h) If $w_1 = i$, $w_2 = i+1$, and $w_3 = i$, then $\Psi = \Phi^{i,i+1,i,w_4,\ldots,w_r} = (\Phi^{i,i+1,w_4,\ldots,w_r})\alpha_{(i,\,i+1,\,i,\,i+1,\,i)}$.

(i) If $w_1 = i$, $w_2 = i+1$, and $w_3 = i+2$, then $i \leq n-3$ and $n \geq 4$, and $\Psi = \Phi^{i,i+1,i+2,w_4,\ldots,w_r} = (\Phi^{i,i+1,w_4,\ldots,w_r})\alpha_{(i,\,i+1,\,i+2,\,i+1,\,i)}$.

Since every sequence $(w_1, \ldots, w_r)$ fits into one of the above cases (except for the singleton sequence (i), which cannot arise), the claim follows. $\square$

Let us consider an example. Let $\mathcal{P}$ be a p-gonal prism, with $p \neq 4$. Then $\mathcal{P}$ is a three-orbit polytope in class $3_{1,2}$, and the flags in the black orbit are those that include the lateral edges. By Theorem 5.3, $\Gamma(\mathcal{P}) = \langle \rho_0, \rho_2, \alpha_{101}, \alpha_{12121} \rangle$. Since the vertices of $\mathcal{P}$ are all 3-valent, the automorphism $\rho_2\alpha_{12121}$ sends the base flag Φ to $\Phi^{1,2,1,2,1,2} = \Phi$. Therefore, $\rho_2\alpha_{12121} = \varepsilon$, and thus $\alpha_{12121} = \rho_2^{-1} = \rho_2$, and the automorphism α_{12121} is redundant. Further analysis shows that $(\rho_0\rho_2)^2 = (\rho_0\alpha_{101})^2 = \varepsilon$. The automorphism $\rho_2\alpha_{101}$ rotates the prism by one face counterclockwise, and thus $(\rho_2\alpha_{101})^p = \varepsilon$. Since $\mathcal{P}$ has $4p$ flags in each flag orbit, and the group

$$\langle \rho_0, \rho_2, \alpha_{101} \mid \rho_0^2, \rho_2^2, \alpha_{101}^2, (\rho_0\rho_2)^2, (\rho_0\alpha_{101})^2, (\rho_2\alpha_{101})^p \rangle$$

is isomorphic to the Coxeter group $[2, p]$ of order $4p$, we see that the group above is the whole group $\Gamma(\mathcal{P})$.

6 Conclusions

The orbit graph of a polytope provides a convenient way to classify k-orbit polytopes, and many other polytope properties follow from properties of the orbit graph. We have developed several necessary properties for orbit graphs, and an obvious question is whether these properties are also sufficient. In other words, given an arc-labeled graph that satisfies Proposition 3.1 and Proposition 3.2, is there a polytope with that orbit graph?

We were able to determine generators for the automorphism group of any three-orbit polytope. The arguments used can be adapted for other orbit graphs as well, but the case analysis becomes unwieldy. It would be helpful to have a general framework with which we can more easily express the automorphisms of a k-orbit polytope. Furthermore, we would like to find other nice properties of these automorphism groups, in the same vein as the "intersection conditions" for the automorphism groups of two-orbit polytopes (see [3]). Finally, we would like to have a way to build a k-orbit polytope directly from a group of the appropriate type, as we can with regular polytopes (see [5, Ch. 2E]). Hubard has done this for two-orbit polyhedra in [3], but generalizing the construction used there does not seem entirely straightforward.

References

[1] Antonio Breda D'Azevedo, *A theory of restricted regularity of hypermaps*, J. Korean Math. Soc **43** (2006), no. 5, 991–1018.

[2] Michael I. Hartley, Daniel Pellicer, and Gordon Williams, *Minimal covers of the prisms and antiprisms*, Discrete Math. **312** (2012), no. 20, 3046–3058.

[3] Isabel Hubard, *Two-orbit polyhedra from groups*, Eur. J. Comb. **31** (2010), no. 3, 943–960.

[4] Isabel A. Hubard, *From geometry to groups and back: The study of highly symmetric polytopes*, ProQuest LLC, Ann Arbor, MI, 2007, Thesis (Ph.D.)–York University (Canada). MR 2712832

[5] Peter McMullen and Egon Schulte, *Abstract regular polytopes*, Encyclopedia of Mathematics and its Applications, vol. 92, Cambridge University Press, Cambridge, 2002. MR 1965665 (2004a:52020)

[6] Mark Mixer, Egon Schulte, and Asia Ivić Weiss, *Hereditary polytopes*, arXiv preprint arXiv:1206.1647, 2012.

[7] Barry Monson and Egon Schulte, *Finite polytopes have finite regular covers*, arXiv preprint arXiv:1209.1167, 2012.

[8] Alen Orbanić, Daniel Pellicer, and Asia Ivić Weiss, *Map operations and k-orbit maps*, J. Combin. Theory Ser. A **117** (2010), no. 4, 411–429.

[9] Daniel Pellicer, *Developments and open problems on chiral polytopes*, Ars Mathematica Contemporanea **5** (2012), no. 2, 333–354.

[10] Egon Schulte and Asia Ivić Weiss, *Chiral polytopes*, Applied geometry and discrete mathematics, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 4, Amer. Math. Soc., Providence, RI, 1991, pp. 493–516. MR 1116373 (92f:51018)